


Paracetamol overdose presenting >24hrs

(Ingested total overdose in ≤ 1 hour time period)



*Clinical judgement required

- Bedside hepatic toxicity: Jaundice, tender liver, hypoglycaemia, encephalopathy, unexplained lactic acidosis.
- Ensure no doubt about time of ingestion or type.
- If uncertainty then treat and review with bloods.

**Clinical judgement required


- Some patients have a chronically raised ALT/INR.
- Review old LFTs/INRs and if chronic derangement discuss with a senior clinician before proceeding to NAC.

Blood monitoring

- Checking a paracetamol level 2hrs before the end of bag 2 is NEW for this protocol.
- U&E, HCO₃, glu, LFTs, FBC and INR should be done 2hrs before the end of each infusion 2. Ensure results are READY for the end of the infusion.
- If unable to achieve blood sampling at the correct time and a delay of >90 minutes is predicted then proceed to the next infusion to avoid prolonged omission of NAC. Bloods should be checked at the earliest opportunity and discontinuation criteria referred to.
- Capillary Blood Glucose (CBG) 6 hourly while on NAC.
- If rapid or progressive biochemical deterioration then discuss with senior and consider referral to regional transplant centre.
- IV NAC can be associated with minor rise in INR without an acute liver injury. (see guideline for details)

*** Bloods should be done 2 hours before the end of infusion 3 and 4.

Paracetamol treatment nomogram and 12-hr shortened N-acetylcysteine dosing schedule (SNAP protocol)



Reproduced courtesy of MHRA

If unclear which of the five paracetamol overdose protocols to follow, discuss with ED / paed's Reg / Cons.

In situations where paracetamol levels will be used to determine need for acetylcysteine (refer to appropriate protocol), plot the measured plasma concentration (in mg/L) against the time since ingestion. If plasma level falls above the line then give acetylcysteine as detailed below.

The nomogram is less accurate between 15-24 hours and accurate ingestion time is even more vital.

Actual weight should be used for calculating both the toxic dose and the acetylcysteine dose - up to a **maximum of 110 kg**

Reactions to acetylcysteine include flushing, nausea & vomiting. Please use 'Acetylcysteine Antidote Adverse Effects – Features & Management' guidance to document any adverse events and guide further management.

Hypersensitivity and anaphylactoid reactions with acetylcysteine are not contraindications as the benefit of treatment still outweighs the risk of not treating.

True anaphylaxis is rare with acetylcysteine but can be managed by stopping the infusion and then restarting at a slower rate.

Table 1. 12-hr shortened N-acetylcysteine dosing schedule (SNAP protocol).

Regimen	First infusion		Second (& extended) infusion	
Infusion fluid	200mL sodium chloride 0.9% or 5% glucose		1000mL sodium chloride 0.9% or 5% glucose	
Preparation	Use 250mL infusion bag and remove 50mL and add required volume of acetylcysteine		Add required volume of acetylcysteine to 1000mL infusion bag	
Duration of infusion	2 hours		10 hours	
Drug dose	100mg/kg acetylcysteine		200mg/kg acetylcysteine	
Weight (kg)	Ampoule volume (mL)	Infusion rate (mL/h)	Ampoule volume (mL)	Infusion rate (mL/h)
30-39	18	109	35	104
40-49	23	112	45	105
50-59	28	114	55	106
60-69	33	117	65	107
70-79	38	119	75	108
80-89	43	122	85	109
90-99	48	124	95	110
100-109	53	127	105	111
≥ 110	55	128	110	111

Each ampoule = 200mg/mL acetylcysteine. Dose calculation based on weight in middle of band. Ampoule rounded up to nearest whole number.